

Impact of 2014/2015 influenza season in Alentejo Region of Portugal

Eleonora Paixão, Paula Valente, Ana Mendes, Sandra Guerra, Anabela David, José Marques Robalo
Alentejo Regional Administration of Health



BACKGROUND

Influenza activity in Europe started week 50/2014 with subtype A(H3N2) viruses as dominant (Broberg *et al.*, 2015). In Portugal was reported medium intensity of influenza activity in week 1/2015 to European Centre for Disease Prevention and Control, with B as dominant subtype.

Influenza causes respiratory infections in humans, leads to increased consultations in general practice and is an important global cause of hospital admissions and mortality.

OBJECTIVE

The main purpose of this study is the estimation of impact of the 2014/2015 influenza season in excess mortality, consultations in primary healthcare, hospital admissions and emergency episodes in Alentejo, Portugal.




Figure 1. Portugal and Alentejo Region

MATERIAL AND METHODS

Mortality data were obtained from daily mortality surveillance system (VDM) provided by National Institute of Health. The number of consultations by flu syndrome (code R80 in International Classification of Primary Care, ICPC-2) and vaccines administered were obtained from SIARS - Informatic System of Regionals Administrations of Health. The number of hospital admissions and emergency episodes were obtained from all five local hospitals in ALERT ADW and SONHO informatics systems.

The period studied, observed period (O), was week 40 to 20 (Monday to Sunday) 2014/2015 season. Expected number (E) was obtained by taking the sum of variables from an equivalent set of reference with same days of week and month from 2013/2014 season. Only one season was considered for reference period, because for some variables other seasons data wasn't available. Excess was calculated for each variable as O-E.

Rate ratios (RR), $RR=O/E$, comparing observed versus reference period, were calculated per variable and week. Confidence intervals (CI) at 95% level used methodology described by Hoshinko *et al.* (2009) and founded by Rothman and Greenland (1998):

$$e^{\left(\ln(RR) \pm 1,96 \cdot \sqrt{\frac{1}{O} + \frac{1}{E}}\right)}$$

where 1,96, is the quantil of normal distribution.

RESULTS




Figure 2. Influenza vaccines administered in Alentejo per week in 2014/2015 season




Figure 3. Estimate of incidence rate, calculated through number of consultations by flu (R80) in primary health care, from 2009/2010 to 2014/2015 season in Alentejo.




Figure 4. RR and 95% CI of consultations by flu (R80) in primary health care from week 40 to 20, 2014/2015 season in Alentejo.




Figure 5. RR and 95% CI of hospital admissions from week 40 to 20, 2014/2015 season in Alentejo.




Figure 6. RR and 95% CI of emergency episodes by flu from week 40 to 20, 2014/2015 season in Alentejo.




Figure 7. RR and 95% CI of total of emergency pediatric episodes from week 40 to 20, 2014/2015 season in Alentejo.




Figure 8. RR and 95% CI of mortality from week 40 to 20, 2014/2015 season in Alentejo.

Vaccines are free of charge within National Health Service for people with 65 or more years old. 82% of the free vaccines (61.577) were administered during the 1st month, October (figure 2).

The peak of incidence rate occurred in week 5/2015 was 255,5/10⁵ inhabitants. This value exceeded the maximum values in Alentejo 2009 and in Portugal 2012 (figure 3).

The total excess of consultations by R80, in primary health care, in season 2014/2015, was 4.530 and statistical significant ($RR=1,72$; $95\%CI: 1,67-1,78$). RR was statistical significant from week 1 to week 12 (figure 4). Particularly, in week 5/2015 there was an excess of 840 consultations, almost the triple than in the reference period ($RR=2,95$ $95\%CI: 2,65-3,30$).

Hospital admissions didn't show a significant excess during the epidemic weeks (figure 5). Total excess was 808 ($RR=1,00$; $95\%CI: 0,99-1,02$).

Week 1 to 14/2015, showed a significant excess of emergency episodes by flu (figure 6). The excess was 1.387 episodes, with statistical significant ($RR=2,08$; $95\%CI: 1,95-2,23$). Week 6/2015 registered the peak of excess (205) of emergency episodes by flu ($RR=5,76$; $95\%CI: 4,17-7,97$).

Total of pediatric emergency showed an excess of 2.916 episodes comparing with reference period, and it was statistical significant with a $RR=1,01$ ($95\%CI: 1,04-1,07$). Week 3/2015, with an excess of 358, $RR=1,29$ ($95\%CI: 1,19-1,38$), was statistical significant (figure 7).

Total excess of mortality was 669, $RR=1,07$ ($95\%CI: 1,03-1,11$). The peak, also in week 3/2015, had 118 excess deaths, comparing with previous season $RR=1,61$ ($95\%CI: 1,35-1,93$), statistical significant.

DISCUSSION/ CONCLUSION

The peak of influenza activity, in Alentejo, was reached in week 5/2015, with an incidence rate, estimated by number of consultations (R80, ICPC-2) in primary health care of 255,5/100.000 inhabitants.

These results are obtained after the end of influenza season. Specific flu mortality data, hospital admissions, and pediatric emergency episodes and other respiratory infections are not yet available. Codification of causes is being done.

The excess mortality can be overestimated, because the VDM includes municipalities that aren't in influence area of Alentejo Regional Administration of Health.

Consultations by flu in primary health care, can be underestimated, because it depends of registries made by general practitioners.

The early detection of influenza epidemics, will allow better planning of the available resources. So in the near future we pretend to establish epidemic thresholds by Moving Epidemic Method (Vega *et al.*, 2012) since this method provides a robust and specific signal to detect influenza epidemics. On the other hand, to estimate disease burden attributable to influenza, other statistical methods, like Poisson regression models, can be used also in future (Schanzer *et al.*, 2013).

Flu surveillance in primary and secondary healthcare should continue, be enhanced or implemented to facilitate early public health risk assessment, prepare health services as well management and treatment of severe cases.

REFERENCES

- Broberg E. *et al.* (2015) Start of the 2014/15 influenza season in Europe: drifted influenza A(H3N2) viruses circulate as dominant subtype.
- Hoshiko S, English P, Smith D, Trent R (2009) A simple method for estimating excess mortality due to heat waves, as applied to the 2006 California heat wave, Int J Public Health, published online.
- Rothman KJ, Greenland S (1998) Modern Epidemiology, 2nd edn. Lippencott-Raven, Philadelphia.
- Schanzer DL, Sevenhuysen C, Winchester B, (2013) Estimating Influenza deaths in Canada, 1992-2009, PLoS ONE 8 (11): e80481.
- Vega T *et al.* (2012) Influenza surveillance in Europe: establishing epidemic thresholds by the Moving Epidemic Method. Influenza and Other Respiratory Viruses 7 (4), 546-558.

ACKNOWLEDGEMENTS

The authors thanks to the Administration Councils of the Local Health Units, the Administration Councils of Évora Hospital and particularly to Ana Coelho, Ana Dionísio, Hugo Quintino, Inês Parrinha, Maria Teresa Mouraia, Sérgio Cardoso, Vasco Carvalho and Vítor Fialho for supplying data.